

FUNCTIONAL EQUATIONS FOR ZETA FUNCTIONS OF \mathbb{F}_1 -SCHEMES

OLIVER LORSCHIED

ABSTRACT. For a scheme X whose \mathbb{F}_q -rational points are counted by a polynomial $N(q) = \sum a_i q^i$, the \mathbb{F}_1 -zeta function is defined as $\zeta_{\mathcal{X}}(s) = \prod (s - i)^{-a_i}$. Define $\chi = N(1)$. In this paper we show that if X is a smooth projective scheme, then its \mathbb{F}_1 -zeta function satisfies the functional equation $\zeta_{\mathcal{X}}(n - s) = (-1)^{\chi} \zeta_{\mathcal{X}}(s)$. We further show that the \mathbb{F}_1 -zeta function $\zeta_G(s)$ of a split reductive group scheme G of rank r with N positive roots satisfies the functional equation $\zeta_G(r + N - s) = (-1)^{\chi} (\zeta_G(s))^{(-1)^r}$.

1. INTRODUCTION

In recent years around a dozen different suggestion of what a scheme over \mathbb{F}_1 should be appeared in literature (cf. [6]). The common motivation for all these approaches is to provide a framework in which Deligne's proof of the Weyl conjectures can be transferred to characteristic 0 in order to proof the Riemann hypothesis. Roughly speaking, \mathbb{F}_1 should be thought of as a field of coefficients for \mathbb{Z} , and \mathbb{F}_1 -schemes \mathcal{X} should have a base extension $\mathcal{X}_{\mathbb{Z}}$ to \mathbb{Z} which is a scheme in the usual sense.

Though it is not clear yet whether one of the existing \mathbb{F}_1 -geometries comes close to this goal, and thus in particular it is not clear what the appropriate notion of an \mathbb{F}_1 -scheme should be, the zeta function $\zeta_{\mathcal{X}}(s)$ of such an elusive \mathbb{F}_1 -scheme \mathcal{X} is determined by the scheme $X = \mathcal{X}_{\mathbb{Z}}$.

Namely, let X be a variety of dimension n over \mathbb{Z} , i.e. a scheme such that X_k is an variety of dimension n for any field k . Assume further that X has a counting polynomial

$$N(q) = \sum_{i=0}^n a_i q^i \in \mathbb{Z}[q],$$

i.e. the number of \mathbb{F}_q -rational points is counted by $\#X(\mathbb{F}_q) = N(q)$ for every prime power q . If X descends to an \mathbb{F}_1 -scheme \mathcal{X} , i.e. $\mathcal{X}_{\mathbb{Z}} \simeq X$, then \mathcal{X} has the zeta function

$$\zeta_{\mathcal{X}}(s) = \lim_{q \rightarrow 1} (q - 1)^{\chi} \zeta_X(q, s)$$

where $\zeta_X(q, s) = \exp\left(\sum_{r \geq 1} N(q^r) q^{-sr} / r\right)$ is the zeta function of $X \otimes \mathbb{F}_q$ if q is a prime power and $\chi = N(1)$ is the order the pole of $\zeta_X(q, s)$ in $q = 1$ (cf. [9]). This expression comes down to

$$\zeta_{\mathcal{X}}(s) = \prod_{i=0}^n (s - i)^{-a_i}$$

([9, Lemme 1]).

From this it is clear that $\zeta_{\mathcal{X}}(s)$ is a rational function in s and that its zeros (resp. poles) are at $s = i$ of order $-a_i$ for $i = 0, \dots, n$. The only statement from the Weyl conjectures which is not obvious for zeta functions of \mathbb{F}_1 -schemes is the functional equation.

I like to thank Takashi Ono for drawing my attention to the symmetries occurring in the counting polynomials of split reductive group schemes, and I like to thank Markus Reineke for his explanations on the comparison theorem for liftable smooth varieties.

2. THE FUNCTIONAL EQUATION FOR SMOOTH PROJECTIVE \mathbb{F}_1 -SCHEMES

Let X be an (irreducible) smooth projective variety of dimension n with a counting polynomial $N(q)$. Let b_0, \dots, b_{2n} be the Betti numbers of X , i.e. the dimensions of the singular homology groups $H_0(X_{\mathbb{C}}), \dots, H_{2n}(X_{\mathbb{C}})$. By Poincaré duality, we know that $b_{2n-i} = b_i$. As a consequence of the comparison theorem for smooth liftable varieties and Deligne's proof of the Weil conjectures, we know that the counting polynomial is of the form

$$N(q) = \sum_{i=0}^n b_{2i} q^i$$

and that $b_i = 0$ if i is odd (cf. [2] and [8]). Thus $\chi = \sum_{i=0}^n b_{2i}$ is the Euler characteristic of $X_{\mathbb{C}}$ in this case (cf. [4]).

Suppose X has an elusive model \mathcal{X} over \mathbb{F}_1 . Then \mathcal{X} has the zeta function $\zeta_{\mathcal{X}}(s) = \prod_{i=0}^n (s - i)^{-b_{2i}}$.

Theorem 1. *The zeta function $\zeta_{\mathcal{X}}(s)$ satisfies the functional equation*

$$\zeta_{\mathcal{X}}(n - s) = (-1)^{\chi} \zeta_{\mathcal{X}}(s)$$

and the factor equals -1 if and only if n is even and b_n is odd.

Proof. We calculate

$$\begin{aligned} \zeta_{\mathcal{X}}(n - s) &= \prod_{i=0}^n ((n - s) - i)^{-b_{2i}} \\ &= \prod_{i=0}^n (-1)^{b_{2i}} (s - (n - i))^{-b_{2i}} \\ &= (-1)^{\chi} \prod_{i=0}^d (s - (n - i))^{-b_{2n-2i}} \end{aligned}$$

where we used $b_{2n-2i} = b_{2i}$ in the last equation. If we now substitute i by $n - i$ in this expression, we obtain

$$\zeta_{\mathcal{X}}(n - s) = (-1)^{\chi} \prod_{i=0}^n (s - i)^{-b_{2i}} = (-1)^{\chi} \zeta_{\mathcal{X}}(s).$$

If n is odd, then there is an even number of non-trivial Betti numbers and $\chi = 2b_0 + 2b_2 + \dots + 2b_{n-1}$ is even. If n is even, then $\chi = 2b_0 + 2b_2 + \dots + 2b_{n-2} + b_n$ has the same parity as b_n . Thus the additional statement. \square

Remark. Note the similarity with the functional equation for motivic zeta functions as in [3, Thm. 1]. Amongst other factors, also $(-1)^{\chi(M)}$ appears in the functional equation of the zeta function of a motive M where $\chi(M)$ is the (positive part of the) Euler characteristic of M .

3. THE FUNCTIONAL EQUATION FOR REDUCTIVE GROUPS OVER \mathbb{F}_1

The above observations imply further a functional equation for reductive group schemes over \mathbb{F}_1 . Note that Soule's and Connes and Consani's approaches towards \mathbb{F}_1 -geometry indeed succeeded in descending split reductive group schemes from \mathbb{Z} to \mathbb{F}_1 (cf. [1], [5], [7]).

Let G be a split reductive group scheme of rank r with Borel group B and maximal split torus $T \subset B$. Let N be the normalizer of T in G and $W = N(\mathbb{Z})/T(\mathbb{Z})$ be the Weyl group. The Bruhat decomposition of G (with respect to T and B) is the morphism

$$\coprod_{w \in W} BwB \longrightarrow G,$$

induced by the subscheme inclusions $BwB \rightarrow G$, which has the property that it induces a bijection between the k -rational points for every field k . We have $B \simeq \mathbb{G}_m^r \times \mathbb{A}^N$ as schemes where N is the number of positive roots of G , and $BwB \simeq \mathbb{G}_m^r \times \mathbb{A}^{N+\lambda(w)}$ where $\lambda(w)$ is the length of $w \in W$. With this we can calculate the counting polynomial of G as

$$N(q) = \# \coprod_{w \in W} BwB(\mathbb{F}_q) = (q-1)^r q^N \sum_{w \in W} q^{\lambda(w)}.$$

The quotient variety G/B is a smooth projective scheme of dimension N with counting function $N_{G/B}(q) = ((q-1)^r q^N)^{-1} N(q) = \sum_{w \in W} q^{\lambda(w)}$. Let b_0, \dots, b_{2N} be the Betti numbers of G/B , then we know from the previous section that $N_{G/B}(q) = \sum_{l=0}^N b_{2l} q^l$ and that $b_{2N-2l} = b_{2l}$.

Thus we obtain for the counting polynomial of G that

$$\begin{aligned} N(q) &= q^N \left(\sum_{k=0}^r (-1)^{r-k} \binom{r}{k} q^k \right) \cdot \left(\sum_{l=0}^N b_{2l} q^l \right) \\ &= \sum_{i=0}^d \left(\sum_{k+l=i-N} (-1)^{r-k} \binom{r}{k} b_{2l} \right) q^i \end{aligned}$$

where $d = r + 2N$ is the dimension of G and with the convention that $\binom{r}{k} = 0$ if $k < 0$ or $k > r$. Denote by $a_i = \sum_{k+l=i-N} (-1)^{r-k} \binom{r}{k} b_{2l}$ the coefficients of $N(q)$.

Lemma. We have $a_0 = \dots = a_{N-1} = 0$ and $a_{d-i} = (-1)^r a_{i+N}$.

Proof. The first statement follows from the fact that $N(q)$ is divisible by q^N . For the second statement we use the symmetries $\binom{r}{k} = \binom{r}{r-k}$ and $b_{2N-2l} = b_{2l}$ to calculate

$$\begin{aligned} a_{d-i} &= \sum_{k+l=d-i-N} (-1)^{r-k} \binom{r}{k} b_{2l} \\ &= \sum_{k+l=d-i-N} (-1)^r (-1)^k \binom{r}{r-k} b_{2N-2l}. \end{aligned}$$

When we substitute k by $r-k$ and l by $N-l$ in this equation and use $d = r + 2N$, then we obtain

$$a_{d-i} = (-1)^r \sum_{k+l=(i+N)-N} (-1)^{r-k} \binom{r}{k} b_{2l},$$

which is the same as $(-1)^r a_{i+N}$. \square

Suppose G has an elusive model \mathcal{G} over \mathbb{F}_1 . Then \mathcal{G} has the zeta function $\zeta_{\mathcal{G}}(s) = \prod_{i=0}^n (s-i)^{-a_i}$. Let $\chi = N(1) = \sum_{i=0}^d a_i$.

Theorem 2. The zeta function $\zeta_{\mathcal{G}}(s)$ satisfies the functional equation

$$\zeta_{\mathcal{G}}(r+N-s) = (-1)^{\chi} \left(\zeta_{\mathcal{G}}(s) \right)^{(-1)^r}.$$

Proof. We use of the previous lemma and $r + N = d - N$ to calculate that

$$\begin{aligned}\zeta_G(r + N - s) &= \prod_{i=0}^n (r + N - s - i)^{-a_i} \\ &= \prod_{i=0}^n (d - N - s - i)^{-(-1)^r a_{d-N-i}}.\end{aligned}$$

After substituting i by $d - N - i$, we find that

$$\begin{aligned}\zeta_G(d - N - s) &= \prod_{i=0}^n (i - s)^{-(-1)^r a_i} \\ &= (-1)^{\sum a_i} \left(\prod_{i=0}^n (s - i)^{-a_i} \right)^{(-1)^r} \\ &= (-1)^{\chi} \left(\zeta_G(s) \right)^{(-1)^r}.\end{aligned}\quad \square$$

Remark. Kurokawa calculates the \mathbb{F}_1 -zeta functions of \mathbb{P}^n , $\mathrm{GL}(n)$ and $\mathrm{SL}(n)$ in [4]. One can verify the functional equation for these examples immediately.

REFERENCES

- [1] A. Connes and C. Consani. *On the notion of geometry over \mathbb{F}_1* . Preprint, [arXiv:0809.2926](#), 2008.
- [2] A. Deitmar. *Remarks on zeta functions and K-theory over \mathbb{F}_1* . Proc. Japan Acad. Ser. A Math. Sci., vol. 82, 2006.
- [3] B. Kahn. *Motivic zeta functions of motives*. Pure and Applied Mathematics Quarterly 5, 1 (2009) 507–570
- [4] N. Kurokawa. *Zeta functions over \mathbb{F}_1* . Proc. Japan Acad. Ser. A Math. Sci. 81 (2005), no. 10, 180–184.
- [5] J. López Peña and O. Lorscheid. *Torified varieties and their geometries over \mathbb{F}_1* . Online first at Mathematische Zeitschrift, 2009.
- [6] J. López Peña and O. Lorscheid. *Mapping \mathbb{F}_1 -land: an overview over geometries over the field with one element*. To be published in the proceeding of the conferences on \mathbb{F}_1 , [arXiv:0909.0069](#), 2009.
- [7] O. Lorscheid. *Algebraic groups over the field with one element*. [arXiv:0907.3824](#), 2009.
- [8] M. Reineke. *Moduli of representations of quivers*. Trends in representation theory of algebras and related topics, EMS Ser. Congr. Rep. (2008), 589–637.
- [9] C. Soulé. *Les variétés sur le corps à un élément*. Mosc. Math. J. 4 (2004), 217–244.

CITY COLLEGE OF CUNY, DEPT. MATH., 160 CONVENT AVE., NYC, NY 10031, U.S.
E-mail address: olorscheid@ccny.cuny.edu